

PROVA 3

1) La vaccinazione antitetanica: aspetti di sorveglianza sanitaria.

2) Chi può rilasciare il documento di esenzione vaccinale per SARS-Co-V2?

Che differenza c'è tra hardware e software?

Coronavirus disease 2019 (Covid-19) has affected tens of millions of people globally since it was declared a pandemic by the World Health Organization on March 11, 2020. Older adults, persons with certain coexisting conditions, and front-line workers are at highest risk for Covid-19 and its complications. Recent data show increasing rates of severe acute respiratory syndrome coronavirus 2 (SARS-CoV2) infection and Covid-19 in other populations, including younger adults.



PROVA 4

1) Che cos'è un giudizio di idoneità e chi sono i destinatari di esso?

2) Elencare le categorie di soggetti per cui è raccomandata la vaccinazione antinfluenzale.

Cosa deve essere digitato nella cella C5 per sommare i valori delle celle C1, C2, C3 e C4 ? (Excel)



a. =SOMMA(C1+C2+C3+C4)

b. =SOMMA(C1+C4)

c. SOMMA(C1+C4)=

d. =SOMMA(C1:C4)

Adults 16 years of age or older who were healthy or had stable chronic medical conditions, including but not limited to human immunodeficiency virus (HIV), hepatitis B virus, or hepatitis C virus infection, were eligible for participation in the trial. Key exclusion criteria included a medical history of Covid-19, treatment with immune-suppressive therapy, or diagnosis with an immunocompromising condition.

PROVA NON ESTRATTA
 

PROVA 2

1) Elencare le vaccinazioni raccomandate per gli operatori sanitari.

2) Come è corretto gestire un infortunio da taglio in sanità?

Quali sono le modalità di trasferimento dei dati fra diversi PC?

Severe systemic events were reported in less than 2% of vaccine recipients after either dose, except for fatigue (in 3.8%) and headache (in 2.0%) after the second dose. Fever (temperature, $\geq 38^{\circ}\text{C}$) was reported after the second dose by 16% of younger vaccine recipients and by 11% of older recipients. Only 0.2% of vaccine recipients and 0.1% of placebo recipients reported fever (temperature, 38.9 to 40°C) after the first dose, as compared with 0.8% and 0.1%, respectively, after the second dose.



PROVA 1

1) Descrivere le attuali indicazioni di gestione di caso COVID positivo, non vaccinato.

2) Elencare le controindicazioni principali alle vaccinazioni.

Che cos'è un database?

Systemic events were reported more often by younger vaccine recipients (16 to 55 years of age) than by older vaccine recipients (more than 55 years of age) in the reactogenicity subset and more often after dose 2 than dose 1. The most commonly reported systemic events were fatigue and headache (59% and 52%, respectively, after the second dose, among younger vaccine recipients; 51% and 39% among older recipients).

